UDS Table 6B Quality of Care Measures Green= LCHD performed better than the 2023 National average. Yellow= LCHD underperformed the 2023 National average. An * indicates that LCHD improved from 2023 to 2024.	2023 National Average	2023 Illinois Average	LCHD 2019	LCHD 2020	LCHD 2021	LCHD 2022	LCHD 2023	LCHD 2024			
Childhood Immunization Status											
MEASURE: Percentage of children 2 years of age who received age appropriate vaccines by their 2nd birthday	30.2%	29.9%	33.5%	37.5%	36.2%	42.2%	44.5%	38.2%			
Cervical Cancer Screening											
MEASURE: Percentage of women 23-64 years of age who were screened for cervical cancer	55.0%	61.5%	71.8%	68.8%	68.8%	73.5%	69.6%	67.3%			
Breast Cancer Screening											
MEASURE: Percentage of women 51-73 years of age who had a mammogram to screen for breast cancer	52.4%	49.4%	NA	39.6%	50.9%	46.5%	50.2%	50.3%*			
Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents											
MEASURE: Percentage of patients 3–17 years of age with a BMI percentile and counseling on nutrition and	71.5%	80.3%	51.4%	51.2%	59.1%	74.8%	88.1%	90.1%*			
physical activity documented Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan											
MEASURE: Percentage of patients 18 years of age and older with (1) BMI documented and (2) follow-up plan documented if BMI is outside normal parameters	67.1%	72.3%	88.9%	87.5%	91.2%	84.0%	94.4%	90.5%			
Preventive Care and Screening: Tobacco Us	e: Screening a	and Cessation	Intervention	1							
MEASURE: Percentage of patients aged 18 years of age and older who (1) were screened for tobacco use one or more times within 24 months, and (2) if identified to be a tobacco user received cessation counseling	84.9%	89.0%	82.3%	83.6%	84.2%	88.2%	90.2%	89.1%			
Statin Therapy for the Prevention and Treatment of Cardiovascular Disease											
MEASURE: Percentage of patients 21 years of age and older with a diagnosis of ASCVD who were prescribed a statin therapy	77.3%	78.5%	58.1%	63.1%	62.1%	70.7%	71.9%	82.0%*			
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet											
MEASURE: Percentage of patients 18 years of age and older with a diagnosis of IVD or AMI, CABG, or PCI procedure with aspirin or another antiplatelet	75.8%	79.3%	88.1%	82.9%	79.3%	79.3%	86.0%	83.0%			
Colorectal Cance	er Screening										
MEASURE: Percentage of patients 50 through 75 years of age who had appropriate screening for colorectal cancer	41.1%	41.6%	33.5%	33.0%	32.1%	39.3%	44.5%	34.9%			
HIV Linkage	to Care										
MEASURE: Percentage of patients whose first ever HIV diagnosis was made by health center staff within 90 days (30 days in 2020) of that first-ever diagnosis	79.7%	86.5%	100.0%	100.0%	66.7%	100%	100%	100%*			
HIV Screening											
MEASURE: Percentage of patients 15 through 65 years of age who were tested for HIV when within age range	48.5%	59.1%	NA	50.1%	53.6%	60.9%	61.9%	72.4%*			
Preventive Care and Screening: Screening for Depression and Follow-Up Plan											

MEASURE: Percentage of patients 12 years of age and older who were (1) screened for depression with a standardized tool and, if screening was positive, (2) had a follow-up plan documented	71.6%	80.3%	73.4%	75.0%	84.8%	93.9%	92.5%	89.6%			
Depression Remission at Twelve Months											
MEASURE: Percentage of patients 12 years of age and older with major depression or dysthymia who reached remission 12 months (+/- 60 days) after an index event	13.6%	13.7%	NA	4.8%	5.7%	7.3%	8.7%	6.5%			
Dental Sealants for Children between 6-9 Years											
MEASURE: Percentage of children 6 through 9 years of age at moderate to high risk of caries who received a sealant on a first permanent molar	58.8%	57.3%	75.2%	35.2%	85.8%	95.8%	96.1%	94.4%			
Low Birth Weight											
MEASURE: Percentage of babies of health center prenatal care patients born whose birth weight was below normal (less than 2,500 grams).	8.6%	9.9%	8.2%	7.1%	7.8%	7.7%	8.9%	7.3*%			
Controlled Hypertension											
MEASURE: Percentage of patients 18 through 85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (less than 140/90 mmHg) during the measurement period.	65.7%	64.7%	64.3%	61.0%	56.6%	63.3%	67.5%	71.5%*			
Uncontrolled Diabetes											
MEASURE: Percentage of patients 18 through 75 years of age with diabetes who had hemoglobin A1c (HbA1c) greater than 9.0 percent or patients without a test during the measurement period	28.8%	29.8%	35.7%	34.8%	34.4%	30.9%	33.4%	33.6%			