

UDS Table 6B Quality of Care Measures Green= LCHD performed better than the 2020 National average. Yellow= LCHD underperformed the 2020 National average. An * indicates that LCHD improved from 2021 to 2022.	2020 National Average	2020 Illinois Average	LCHD 2019	LCHD 2020	LCHD 2021	LCHD 2022
Childhood Immunization Status						
MEASURE: Percentage of children 2 years of age who received age appropriate vaccines by their 2nd birthday	40.4%	34.9%	33.5%	37.5%	36.2%	42.2%*
Cervical Cancer Screening						
MEASURE: Percentage of women 23-64 years of age who were screened for cervical cancer	51.0%	55.3%	71.8%	68.8%	68.8%	73.5%*
Breast Cancer Screening						
MEASURE: Percentage of women 51-73 years of age who had a mammogram to screen for breast cancer	45.3%	39.0%	NA	39.6%	50.9%	46.5%
Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents						
MEASURE: Percentage of patients 3-17 years of age with a BMI percentile and counseling on nutrition and physical activity documented	65.1%	71.7%	51.4%	51.2%	59.1%	74.8%*
Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan						
MEASURE: Percentage of patients 18 years of age and older with (1) BMI documented and (2) follow-up plan documented if BMI is outside normal parameters	65.7%	67.2%	88.9%	87.5%	91.2%	84.0%
Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention						
MEASURE: Percentage of patients aged 18 years of age and older who (1) were screened for tobacco use one or more times within 24 months, and (2) if identified to be a tobacco user received cessation counseling intervention	83.4%	83.0%	82.3%	83.6%	84.2%	88.2%*
Statin Therapy for the Prevention and Treatment of Cardiovascular Disease						
MEASURE: Percentage of patients 21 years of age and older with a diagnosis of ASCVD who were prescribed a statin therapy	71.9%	71.8%	58.1%	63.1%	62.1%	70.7%*
Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antiplatelet						
MEASURE: Percentage of patients 18 years of age and older with a diagnosis of IVD or AMI, CABG, or PCI procedure with aspirin or another antiplatelet	78.8%	83.0%	88.1%	82.9%	79.3%	79.3%
Colorectal Cancer Screening						
MEASURE: Percentage of patients 50 through 75 years of age who had appropriate screening for colorectal cancer	40.1%	37.9%	33.5%	33.0%	32.1%	39.3%*
HIV Linkage to Care						
MEASURE: Percentage of patients whose first ever HIV diagnosis was made by health center staff within 90 days (30 days in 2020) of that first-ever diagnosis	81.4%	86.2%	100.0%	100.0%	66.7%	100%*
HIV Screening						
MEASURE: Percentage of patients 15 through 65 years of age who were tested for HIV when within age range	32.3%	40.0%	NA	50.1%	53.6%	60.9%*
Preventive Care and Screening: Screening for Depression and Follow-Up Plan						
MEASURE: Percentage of patients 12 years of age and older who were (1) screened for depression with a standardized tool and, if screening was positive, (2) had a follow-up plan documented	64.2%	70.5%	73.4%	75.0%	84.8%	93.9%*
Depression Remission at Twelve Months						
MEASURE: Percentage of patients 12 years of age and older with major depression or dysthymia who reached remission 12 months (+/- 60 days) after an index event	13.7%	12.7%	NA	4.8%	5.7%	7.3%*
Dental Sealants for Children between 6-9 Years						
MEASURE: Percentage of children 6 through 9 years of age at moderate to high risk of caries who received a sealant on a first permanent molar	48.7%	46.6%	75.2%	35.2%	85.8%	95.8%*
Low Birth Weight						
MEASURE: Percentage of babies of health center prenatal care patients born whose birth weight was below normal (less than 2,500 grams).	8.2%	8.4%	8.2%	7.1%	7.8%	7.7%*
Controlled Hypertension						
MEASURE: Percentage of patients 18 through 85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (less than 140/90 mmHg) during the measurement period.	58.0%	55.8%	64.3%	61.0%	56.6%	63.3%*
Uncontrolled Diabetes						
MEASURE: Percentage of patients 18 through 75 years of age with diabetes who had hemoglobin A1c (HbA1c) greater than 9.0 percent or patients without a test during the measurement period	35.6%	38.5%	35.7%	34.8%	34.4%	30.9%*