UDS Clinical Quality Indicators

Indicator and Description	Compliant Patients	Indicator Population	Compliance Percentage
BMI Screening and Follow-Up Percentage of patients 18 years of age and older with BMI documented and follow up plan documented if BMI is less than 18.5 kg/m ² or greater than 25 kg/m ²	10,217	12,252	HRSA Target: 90%
Cervical Cancer Screening Percentage of female patients aged 23-64 who had cervical cytology performed within the last 3 years or women aged 30-64 who had cervical cytology/HPV co-testing performed within the last 5 years	4,577	6,724	68.1%
Childhood Immunization Status Percentage of patients 2 years of age with all of their age-appropriate vaccines (4 DTaP, 3 IPV, 1 MMR, 3 HiB, 3 Hep B, 1 VZV, 4 PCV, 1 Hep A, 2 or 3 RV, and 2 flu)	255	620	41.1%
Colorectal Cancer Screening Percentage of patients 50–75 years of age who had appropriate screening for colorectal cancer	1,027	4,025	25.5%
Dental Sealants for Children Percentage of patients aged 6-9 years at moderate to high risk of caries who recieved a sealant on a first permanent molar	30	40	7\$.0%
Depression Screening and Plan Percentage of patients aged 12 years and older screened for depression using an age-appropriate screening tool and if positive, a follow-up is documented on the date of the positive screen	6,896	11,165	61.8%
Diabetes (Uncontrolled)* Percentage of patients 18–75 years of age with diabetes who had hemoglobin A1c (HbA1c) greater than 9.0 percent during the measurement period	1,103	2,688	41.0%
			0% 20% 40% 60% 80% 100% Compliance Percentage
* Measures are 'negative', meaning that a smaller perc	entage is better		2019 Percentage 2020 YTD Percentage

UDS Clinical Quality Indicators

Indicator and Description	Compliant Patients	Indicator Population	Compliance Percentage
HIV Linkage to Care Percentage of newly diagnosed HIV patients that received treatment within 90 days of diagnosis	6	7	100% 85.7%
Hypertension (Controlled) Percentage of patients 18–85 years of age who had a diagnosis of hypertension and whose blood pressure (BP) was adequately controlled (less than 140/90 mmHg) during the measurement period.	2,240	4,095	70%
Ischemic Vascular Disease Medication Use Percentage of patients aged 18 years and older with IVD or had an acute myocardial infarction, coronary artery bypass graft, or percutaneous coronary interventions in the previous 12 months and used aspirin or an antiplatelet	232	254	87% 91.3%
Low Birth Weights* Percentage of babies of health center prenatal care patients born whose birth weight was below normal (less than 2,500 grams).	20	275	7% 7.3%
Prenatal Care Early Entry Percentage of prenatal patients with their first prenatal visit during their first trimester	457	646	75%
Statin Therapy for Cardiovascular Disease Percentage of patients aged 21 years and older at high risk of cardiovascular events who were prescribed statin therapy	2,520	3,930	60% 64.1%
Tobacco Screening and Cessation Percentage of patients aged 18 and older who were screened for tobacco use within 24 months and who received cessation counseling intervention if defined as a tobacco user	6,504	7,963	81.7%
Weight Counseling for Ages < 18 years Percentage of patients 3-17 years of age with a BMI percentile and counseling on nutrition and physical activity documented	2,067	4,598	45.0%
			0% 20% 40% 60% 80% 100% Compliance Percentage
* Measures are 'negative', meaning that a smaller percentage is b	petter	2019	Percentage 2020 YTD Percentage