
UDS Clinical Quality Indicators


Indicator and Description	Compliant Patients	Indicator Population
BMI Screening and Follow-Up Percentage of patients 18 years of age and older with BMI documented and follow up plan documented if BMI is less than 18.5 kg/m^2 or greater than 25 kg/m^2	4,664	5,081
Breast Cancer Screening Percentage of female patients 51–73 years of age who had a mammogram to screen for breast cancer	611	1,015
Cervical Cancer Screening Percentage of female patients aged 23-64 who had cervical cytology performed within the last 3 years or women aged 30-64 who had cervical cytology/ HPV co-testing performed within the last 5 years	2,207	3,260
Childhood Immunization Status Percentage of patients 2 years of age with all of their age-appropriate vaccines (4 DTaP, 3 IPV, 1 MMR, 3 HiB, 3 Hep B, 1 VZV, 4 PCV, 1 Hep A, 2 or 3 RV, and 2 flu)	139	347
Colorectal Cancer Screening Percentage of patients 50–75 years of age who had appropriate screening for colorectal cancer	558	1,962
Dental Sealants for Children Percentage of patients aged 6-9 years at moderate to high risk of caries who recieved a sealant on a first permanent molar	17	25
Depression Remission Percentage of patients 12 years of age and older with major depression or dysthymia who reached remission 12 months (+/- 60 days) after an index event	6	272
Depression Screening and Plan Percentage of patients aged 12 years and older screened for depression using an age-appropriate screening tool and if positive, a follow-up is documented on the date of the positive screen	3,230	5,188
Diabetes (Uncontrolled)* Percentage of patients 18–75 years of age with diabetes who had hemoglobin A1c (HbA1c) greater than 9.0 percent or patients without a test during the measurement period	803	1,337



* Measures are 'negative', meaning that a smaller percentage is better

UDS Clinical Quality Indicators

Indicator and Description	Compliant Patients	Indicator Population	Compliance Percentage
HIV Linkage to Care Percentage of newly diagnosed HIV patients that received treatment within 90 days of diagnosis	1	1	100%
HIV Screening Percentage of patients 15-65 years of age who were tested for HIV	10,059	20,341	49.5%
Hypertension (Controlled) Percentage of patients 18–85 years of age who had a diagnosis of hypertension and whose blood pressure (BP) was adequately controlled (less than 140/90 mmHg) during the measurement period.	987	2,062	47.9% / 70%
Ischemic Vascular Disease Medication Use Percentage of patients aged 18 years and older with IVD or an acute myocardial infarction, coronary artery bypass graft, or percutaneous coronary intervention in the previous year and used aspirin or an antiplatelet	99	125	79.2% / 87%
Low Birth Weights* Percentage of babies of health center prenatal care patients born whose birth weight was below normal (less than 2,500 grams).	0	34	0.0% / 7%
Prenatal Care Early Entry Percentage of prenatal patients with their first prenatal visit during their first trimester	232	312	74.4% / 75%
Statin Therapy for Cardiovascular Disease Percentage of patients aged 21 years and older at high risk of cardiovascular events who were prescribed statin therapy	1,218	1,891	64.4% / 60%
Tobacco Screening and Cessation Percentage of patients aged 18 and older who were screened for tobacco use within 24 months and who received cessation counseling intervention if defined as a tobacco user	2,030	2,359	86.1% / 88%
Weight Counseling for Ages < 18 years Percentage of patients 3-17 years of age with a BMI percentile and counseling on nutrition and physical activity documented	707	1,582	44.7% / 80%



* Measures are 'negative', meaning that a smaller percentage is better